arXiv:1610.07408v3 [math.CO] 18 Nov 2016

Congruence classes of triangles in $\mathbb{F}_p^2$

Thang Pham Le Anh Vinh

Abstract

In this short note, we give a lower bound on the number of congruence classes of triangles in a small set of points in $\mathbb{F}_p^2$. More precisely, for $\mathcal{A} \subset \mathbb{F}_p^2$ with $|\mathcal{A}| \leq p^{2/3}$, we prove that the number of congruence classes of triangles determined by points in $\mathcal{A} \times \mathcal{A}$ is at least $|\mathcal{A}|^{7/2}$. This note is not intended for journal publication.

1 Introduction

Let $\mathbb{F}_q$ be a finite field of order q with $q = p^r$. Two k-simplices in $\mathbb{F}_q^2$ with vertices $(\mathbf{x}_1, \ldots, \mathbf{x}_{k+1})$, $(\mathbf{y}_1, \ldots, \mathbf{y}_{k+1})$ are in the same congruence class if there exist an orthogonal matrix θ and a vector $\mathbf{z} \in \mathbb{F}_q^d$ such that $\mathbf{z} + \theta(\mathbf{x}_i) = \mathbf{y}_i$ for all $i = 1, 2, \ldots, k+1$. Let $T_{k,d}(\mathcal{E})$ denote the set of congruence classes of k-simplices determined by points in $\mathcal{E}$. In this paper, we use the notation $X \ll Y$ which means that there exists a positive absolute constant C that does not depend on X, Y and q such that $X \leq CY$.

The first result on the size of $T_{k,d}(\mathcal{E})$ in $\mathbb{F}_q^d$ was established by Hart and Iosevich [4]. They indicated that if $|\mathcal{E}| \gg q^{\frac{kd}{k+1}+\frac{k}{2}}$ with $d \geq \binom{k+1}{2}$, then $|T_{k,d}(\mathcal{E})| \gg q^{\binom{k+1}{2}}$. By employing techniques from spectral graph theory, the second listed author [11] proved that $\mathcal{E}$ also contains a copy of all k-simplices with non-zero edges under the conditions $d \geq 2k$ and $|\mathcal{E}| \gg q^{(d-1)/2+k}$. It follows from the results of Hart and Iosevich [4] and Vinh [11] that we only can get a non-trivial result on the number of congruence classes of triangles when $d \geq 4$. There are several progresses on improving this problem. The result, which we have to mention first, is a result due to Covert et al. [3]. They proved that if $|\mathcal{E}| \gg \rho q^2$, then $|T_{2,2}(\mathcal{E})| \gg \rho q^3$. This means that in order to get a constant factor of all congruence classes of triangles in $\mathbb{F}_p^2$, we need the condition $|\mathcal{E}| \gg q^2$. In 2014, Bennett, Iosevich and Pakianathan [1], using Elekes and Sharir's approach in the Erdős distinct distance problem, improved this result, namely, they showed that the condition $|\mathcal{E}| \geq q^{7/4}$ is enough to get at least cq^3 congruence classes of triangles for some absolute positive constant c. Recently, by using Fourier analytic methods and elementary results from group action theory, Bennett, Hart, Iosevich, Pakianathan, and Rudnev [2] improved the threshold $q^{7/4}$ to $q^{8/5}$. In the case $\mathcal{E}$ is the Cartesian product of sets with different sizes, the authors and Hiep [8] obtained the following improvement.

Theorem 1.1 (Theorem 1.2, [8]). *Let $\mathcal{E} = \mathcal{A}_1 \times \mathcal{A}_2$ be a subset in $\mathbb{F}_q^2$. Suppose that*

$$(\min_{1\le i\le 2} |\mathcal{A}_i|)^{-1}|\mathcal{E}|^{k+1} \gg q^{2k},$$

then for $1 \le k \le 2$, we have

$$|T_{k,2}(\mathcal{E})| \gg q^{\binom{k+1}{2}}.$$

The following is a consequence of Theorem 1.1.

Theorem 1.2 ([8]). *Let $\mathcal{A}, \mathcal{B}$ be subsets in $\mathbb{F}_q$. If $|\mathcal{A}| \ge q^{\frac{1}{2}+\epsilon}$ and $|\mathcal{B}| \ge q^{1-\frac{2\epsilon}{3}}$ for some $\epsilon \ge 0$, then we have*

$$|T_{2,2}(\mathcal{A} \times \mathcal{B})| \gg q^3.$$

We note that in [2], Bennett et al. gave a construction of $|\mathcal{A}| = q^{1/2-\epsilon'}$ and $|\mathcal{B}| = q$, and the number of congruence classes triangles determined by $\mathcal{A} \times \mathcal{B}$ is at most $cq^{3-\epsilon"}$ for $\epsilon" > 0$. On the other hand, it follows from Theorem 1.2 that if $|\mathcal{A}| < q^{1/2}$ then we must have $|\mathcal{B}| > q$ to guarantee that $|T_{2,2}(\mathcal{A} \times \mathcal{B})| \gg q^3$. Hence, the condition of $|\mathcal{A}| \cdot |\mathcal{B}|$ in Theorem 1.2 is sharp up to a factor of $q^{\epsilon/3}$. From the construction in [2], Iosevich [6] conjectured that the right size of a general set $\mathcal{E} \subseteq \mathbb{F}_q^2$ for getting at least cq^3 congruence classes of triangles is around $q^{3/2}$.

The main purpose of this short note is to give a lower bound on the number of congruence classes of triangles in $\mathcal{A} \times \mathcal{A}$ when $\mathcal{A}$ is a small set of points.

Theorem 1.3. *Let $\mathbb{F}_p$ be a field of odd prime p. For $\mathcal{A} \subseteq \mathbb{F}_p$ with $|\mathcal{A}| \le p^{2/3}$, we have*

$$|T_{2,2}(\mathcal{A} \times \mathcal{A})| \ge |\mathcal{A}|^{7/2}.$$

2 Two proofs of Theorem 1.3

The first proof: Let $\mathbb{F}_p$ be a prime field of order p. For $\mathcal{E} \subseteq \mathbb{F}_p^2$ and $t \in \mathbb{F}_p$, we define $\nu_{\mathcal{E}}(\lambda)$ as the cardinality of the set $\{(\mathbf{x}, \mathbf{y}) \in \mathcal{E} \times \mathcal{E} \colon ||\mathbf{x} - \mathbf{y}|| = \lambda\}$. In order to prove Theorem 1.3, we first need the following lemmas.

Lemma 2.1 ([8]). *Let $\mathcal{E}$ be a subset in $\mathbb{F}_p^2$. For a fixed $\lambda \in \mathbb{F}_p$, denote by $H_\lambda(\mathcal{E})$ the number of hinges of the form $(\mathbf{p}, \mathbf{q_1}, \mathbf{q_2}) \in \mathcal{E} \times \mathcal{E} \times \mathcal{E}$ with $||\mathbf{p} - \mathbf{q_1}|| = ||\mathbf{p} - \mathbf{q_2}|| = \lambda$. We have the following estimate*

$$\sum_{\lambda \in \mathbb{F}_p} \nu_{\mathcal{E}}(\lambda)^2 \ll \frac{|\mathcal{E}|}{4} \sum_{\lambda \in \mathbb{F}_p} H_\lambda(\mathcal{E}) + |\mathcal{E}|^3. \tag{2.1}$$

By applying the recent Rudnev's breakthrough on the number of incidences between points and planes in $\mathbb{F}_p^3$ in [9], Petridis [7] obtained the following.

Lemma 2.2 (**Petridis**, [7]). *Let $\mathbb{F}_p$ be a field of order odd prime p. For $\mathcal{A} \subseteq \mathbb{F}_p$ with $|\mathcal{A}| \le p^{2/3}$, we have*

$$\sum_{\lambda \in \mathbb{F}_p} H_\lambda(\mathcal{A} \times \mathcal{A}) \ll |\mathcal{A}|^{9/2}.$$

As a consequence of Theorem 2.2, Petridis [7] proved the following result on the number of distinct distances in $\mathcal{A} \times \mathcal{A}$.

Theorem 2.3 (**Petridis**, [7]). *Let $\mathbb{F}_p$ be a field of order odd prime p. For $\mathcal{A} \subseteq \mathbb{F}_p$ with $|\mathcal{A}| \leq p^{2/3}$, we have the number of distinct distances determined by points in $\mathcal{A} \times \mathcal{A}$ is at least $\min\{p, |\mathcal{A}|^{3/2}\}$.*

Combining Lemma 2.2 and Lemma 2.1, we obtain the following.

Lemma 2.4. *Let $\mathbb{F}_p$ be a field of order odd prime p. For $\mathcal{A} \subseteq \mathbb{F}_p$ with $|\mathcal{A}| \leq p^{2/3}$, we have*

$$\sum_{\lambda \in \mathbb{F}_p} \nu_{\mathcal{A}\times\mathcal{A}}(\lambda)^2 \ll |A|^{13/2}. \tag{2.2}$$

Proof of Theorem 1.3. On one hand, by applying the Cauchy-Schwarz inequality, we have

$$|T_{2,2}(\mathcal{A} \times \mathcal{A})| \geq \frac{|\mathcal{A}|^{12}}{N},$$

where N is the number of pairs of congruence triangles. On the other hand, two triangles, which are denoted by $\Delta(\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3)$ and $\Delta(\mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3)$, are in the same congruence class if there exist an orthogonal matrix M and a vector $\mathbf{z} \in \mathbb{F}_p^2$ such that

$$M\mathbf{a}_i + \mathbf{z} = \mathbf{b}_i, \ 1 \leq i \leq 3.$$

This implies that

$$N \leq |\mathcal{A} \times \mathcal{A}| \sum_{\lambda \in \mathbb{F}_p} \nu_{\mathcal{A}\times\mathcal{A}}(\lambda)^2.$$

Thus, it follows from Lemma 2.4 that

$$N \leq |\mathcal{A} \times \mathcal{A}| \sum_{\lambda \in \mathbb{F}_p} \nu_{\mathcal{A}\times\mathcal{A}}(\lambda)^2 = |\mathcal{A}|^{17/2}.$$

In other words,

$$|T_{2,2}(\mathcal{A} \times \mathcal{A})| \geq |\mathcal{A}|^{7/2},$$

and the theorem follows. □

The second proof: The following lemma was suggested by Frank de Zeeuw.

Lemma 2.5. *Let $\mathbb{F}_p$ be a field of order odd prime p. For $\mathcal{A} \subseteq \mathbb{F}_p$, we have*

$$|T_{2,2}(\mathcal{A} \times \mathcal{A})| \geq \frac{1}{6}(|\mathcal{A}|^2 - 2) \cdot |\Delta_{\mathbb{F}_p}(\mathcal{A} \times \mathcal{A})|,$$

where $\Delta_{\mathbb{F}_p}(\mathcal{A} \times \mathcal{A})$ is the set of distinct distances determined by points in $\mathcal{A} \times \mathcal{A}$.

Proof. Suppose that $m = |\Delta_{\mathbb{F}_p}(\mathcal{A} \times \mathcal{A})|$, then there exist m segments with distinct distances. For each segment, it is easy to check that there are at least $(|\mathcal{A}|^2-2)/2$ congruence classes of triangles with the same base. On the other hand, when we count congruence classes of triangles with the bases from the set of m segments, each class will be repeated at most 3 times. This implies that

$$|T_{2,2}(\mathcal{A} \times \mathcal{A})| \geq \frac{1}{6}(|\mathcal{A}|^2 - 2) \cdot |\Delta_{\mathbb{F}_p}(\mathcal{A} \times \mathcal{A})|,$$

which concludes the proof of the lemma. □

From Theorem 2.3 and Lemma 2.5, we are able to obtain a slight weaker version of Theorem 1.3 as follows.

Theorem 2.6. *Let $\mathbb{F}_p$ be a field of odd prime p. For $\mathcal{A} \subseteq \mathbb{F}_p$ with $|\mathcal{A}| \leq p^{2/3}$, we have*

$$|T_{2,2}(\mathcal{A} \times \mathcal{A})| \geq \frac{1}{6}|\mathcal{A}|^{7/2}.$$

Let $\mathbb{F}_q$ be a finite field of order q. For $\mathcal{E} \subseteq \mathbb{F}_q^2$ with $|\mathcal{E}| \geq \rho q^2$ with $q^{-1/2} \ll \rho < 1$. Iosevich and Rudnev [5] proved that the number of distinct distances determined by points in $\mathcal{E}$ is at least $q-1$. It follows from Lemma 2.5 that the number of congruence classes of triangles in $\mathcal{E}$ is at least $\rho(q-1)q^2$. In other words, we have recovered the result due to Covert et al. [3] and Vinh [11] for the case $d = 2$.

Theorem 2.7 (**Covert et al.**, [3])**.** *Let $\mathbb{F}_q$ be a finite field of order q. For $\mathcal{E} \subseteq \mathbb{F}_q^2$ with $|\mathcal{E}| \geq \rho q^2$ with $q^{-1/2} \ll \rho < 1$, then the number of congruence classes of triangles in $\mathcal{E}$ is at least $c\rho q^3$, for some positive constant c.*

References

[1] M. Bennett, A. Iosevich, J. Pakianathan, *Three-point configurations determined by subsets of $\mathbb{F}_q^2$ via the Elekes-Sharir Paradigm*, Combinatorica, **34**(6) (2014), 689–706.

[2] M. Bennett, D. Hart, A. Iosevich, J. Pakianathan, M. Rudnev, *Group actions and geometric combinatorics in $\mathbb{F}_q^d$*, to appear in Forum Math., 2016.

[3] D. Covert, D. Hart, A. Iosevich, S. Senger, I. Uriarte-Tuero, *A FurstenbergKatznelsonWeiss type theorem on $(d+1)$-point configurations in sets of positive density in finite field geometries*, Discrete Mathematics, **311**(6) (2011), 423–430.

[4] D. Hart, A. Iosevich, *Ubiquity of simplices in subsets of vector spaces over finite fields*, Analysis Mathematika, **34** (2007).

[5] A. Iosevich, M. Rudnev, *Erdős distance problem in vector spaces over finite fields*, Trans. Am. Math. Soc. **359** (2007), 6127–6142.

[6] A. Iosevich, *Personal communication*, 2016.

[7] G. Petridis, *Pinned algebraic distances determined by Cartesian products in* $\mathbb{F}_p^2$, arxiv: 1610.03172v1.

[8] H. Pham, T. Pham, L. A. Vinh, *An improvement on the number of simplices in* $\mathbb{F}_q^d$, submitted, arxiv: 1608.06398v1.

[9] M. Rudnev, *On the number of incidences between planes and points in three dimensions*, to appear in Combinatorica (2016).

[10] L. A. Vinh, *Triangles in vector spaces over finite fields*, Online Journal of Analytic Combinatorics **6** (2009) 1–8.

[11] L. A. Vinh, *On kaleidoscopic pseudo-randomness of finite Euclidean graphs*, Discussiones Mathematicae: Graph Theory, **32**(2).

Department of Mathematics,
EPF Lausanne
Switzerland
E-mail: thang.pham@epfl.ch

University of Education,
Vietnam National University
Viet Nam
E-mail: vinhla@vnu.edu.vn